

Wound Dressings During Radiotherapy for Cancer: A Survey of Practice

Siby Elizabeth J. Thomas, MSN, RN, Sheryl Reimer-Kirkham, PhD, RN, and Rosemary Kohr, PhD, RN



© Top Photo Group/Thinkstock

Background: Patients undergoing radiotherapy may experience changes to the skin that require dressings. Recommendations regarding radiating through wound dressings have been variable and relate to the concern regarding surface dose increase or bolus effect.

Objectives: The purpose of this article is to identify current evidence and practice through literature review and a national environmental scan.

Methods: Nurses from 18 radiation oncology centers in Canada were surveyed about current practice. In-depth telephone interviews were conducted with four nurse participants to further understand the context of this issue within the nursing practice environment.

Findings: The integrated results of the study were reviewed with five clinical experts to make recommendations for research, practice, leadership, and policy. Implications for clinical practice included the involvement of radiation oncology nurses in the treatment planning team, development of clinical practice tools, and the relevance of the Person-Centered Nursing framework for wound management during radiotherapy.

Siby Elizabeth J. Thomas, MSN, RN, is an education resource nurse at the BC Cancer Agency in Fraser Valley and Abbotsford Centres, Sheryl Reimer-Kirkham, PhD, RN, is a professor in the School of Nursing at Trinity Western University in Langley, both in British Columbia, Canada; and Rosemary Kohr, PhD, RN, is a consultant at Kohr Consulting in London, Ontario, Canada. The authors take full responsibility for the content of the article. This study was supported, in part, by funding from the Fraser Valley Radiation Therapy Research and Innovation Fund Committee and the British Columbia Nurses Union. The content of this article has been reviewed by independent peer reviewers to ensure that it is balanced, objective, and free from commercial bias. No financial relationships relevant to the content of this article have been disclosed by the independent peer reviewers or editorial staff. Description of products does not indicate or imply endorsement by the *Clinical Journal of Oncology Nursing* or the Oncology Nursing Society. Thomas can be reached at sthomas4@bccancer.bc.ca, with copy to editor at CJONEditor@ons.org. (Submitted May 2014. Revision submitted October 2014. Accepted for publication October 14, 2014.)

Key words: wound management; radiotherapy; clinical decision-making; interprofessional practice; person-centered nursing

Digital Object Identifier: 10.1188/15.CJON.E87-E91

Wounds are common among patients with cancer undergoing radiotherapy. The management of these wounds, such as the standard removal of wound dressings prior to daily fractionated radiotherapy, can be a source of discomfort for patients and a challenge for the nurses involved. At a patient's request and with the approval of the radiation oncologist, very thin dressings have been left in place during the radiation treatment in some settings. However, little evidence exists to inform whether removing the dressing alters the wound bed and affects the healing process or whether the radiation dose changes when delivered through a dressing. As suggested by Hollinworth and Mann (2010), "Further consideration could be given to leaving the product in place and the radiotherapy dose recalculated on an individual patient basis . . . during the initial radiotherapy planning stage" (p. 63). In this article, the authors present the results of a study that examined

current evidence and practice in relation to wound dressings during cancer radiotherapy.

Literature Review

Radiation-induced dermatitis is one of the most common side effects of external beam radiotherapy (Harris et al., 2012) and has the potential to affect an individual's quality of life. Patients with dermatitis may suffer in a variety of ways: changes in body image, physical discomfort such as pain and itching, and difficulty with activities of daily living (McQuestion, 2006). If the resulting wounds require a dressing, clinicians have to choose from a plethora of dressing products. Dressings are chosen taking into account goals of care such as managing exudate, facilitating a moist wound environment, and minimizing pain and bleeding. These products often include antimicrobial dressings—silver being the most common or popular among